



Relations and Their Properties



What is a relation

- Let A and B be sets. A binary relation R is a subset of $A \times B$
- Example
 - Let A be the students in a the CS major
 - $A = \{\text{Alice, Bob, Claire, Dan}\}$
 - Let B be the courses the department offers
 - $B = \{\text{CS101, CS201, CS202}\}$
 - We specify relation $R = A \times B$ as the set that lists all students $a \in A$ enrolled in class $b \in B$
 - $R = \{ (\text{Alice, CS101}), (\text{Bob, CS201}), (\text{Bob, CS202}), (\text{Dan, CS201}), (\text{Dan, CS202}) \}$




More relation examples

- Another relation example:
 - Let A be the cities in the US
 - Let B be the states in the US
 - We define R to mean a is a city in state b
 - Thus, the following are in our relation:
 - (C'ville, VA)
 - (Philadelphia, PA)
 - (Portland, MA)
 - (Portland, OR)
 - etc...
- Most relations we will see deal with ordered pairs of integers



Representing relations

We can represent relations graphically:



We can represent relations in a table:

	CS101	CS201	CS202
Alice	X		
Bob		X	X
Claire			
Dan		X	X




Relations on a set

- A relation on the set A is a relation from A to A
 - In other words, the domain and co-domain are the same set
 - We will generally be studying relations of this type



Relations on a set

- Let A be the set $\{ 1, 2, 3, 4 \}$
- Which ordered pairs are in the relation $R = \{ (a,b) \mid a \text{ divides } b \}$
- $R = \{ (1,1), (1,2), (1,3), (1,4), (2,2), (2,4), (3,3), (4,4) \}$



R	1	2	3	4
1	X	X	X	X
2		X		X
3			X	
4				X



More examples

- Consider some relations on the set \mathbf{Z}
- Are the following ordered pairs in the relation?

(1,1) (1,2) (2,1) (1,-1) (2,2)

- $R_1 = \{ (a,b) \mid a \leq b \}$

- $R_2 = \{ (a,b) \mid a > b \}$

- $R_3 = \{ (a,b) \mid a = |b| \}$

- $R_4 = \{ (a,b) \mid a = b \}$

- $R_5 = \{ (a,b) \mid a = b + 1 \}$

- $R_6 = \{ (a,b) \mid a + b \leq 3 \}$

	X	X			X
			X	X	
	X			X	X
	X				X
			X		
	X	X	X	X	



Relation properties

- Six properties of relations we will study:
 - انعكاسية Reflexive
 - لانعكاسية Irreflexive
 - متماثلة Symmetric
 - غير متماثلة Asymmetric
 - ضد متماثلة Antisymmetric
 - متعدية Transitive



Reflexivity

- A relation is reflexive if every element is related to itself
 - Or, $(a,a) \in R$
- Examples of reflexive relations:
 - $=, \leq, \geq$
- Examples of relations that are not reflexive:
 - $<, >$



Irreflexivity

- A relation is irreflexive if every element is *not* related to itself
 - Or, $(a,a) \notin R$
 - Irreflexivity is the opposite of reflexivity
- Examples of irreflexive relations:
 - $<, >$
- Examples of relations that are not irreflexive:
 - $=, \leq, \geq$



Reflexivity vs. Irreflexivity

- A relation can be neither reflexive nor irreflexive
 - Some elements are related to themselves, others are not
- We will see an example of this later on



Symmetry

- A relation is symmetric if, for every $(a,b) \in R$, then $(b,a) \in R$
- Examples of symmetric relations:
 - $=$, isTwinOf()
- Examples of relations that are not symmetric:
 - $<$, $>$, \leq , \geq



Asymmetry

- A relation is asymmetric if, for every $(a,b) \in R$, then $(b,a) \notin R$
 - Asymmetry is the opposite of symmetry
- Examples of asymmetric relations:
 - $<, >$
- Examples of relations that are not asymmetric:
 - $=, \text{isTwinOf}(), \leq, \geq$



Antisymmetry

- A relation is antisymmetric if, for every $(a,b) \in R$, then $(b,a) \in R$ is true only when $a=b$
 - Antisymmetry is *not* the opposite of symmetry
- Examples of antisymmetric relations:
 - $=, \leq, \geq$
- Examples of relations that are not antisymmetric:
 - $<, >, \text{isTwinOf}()$



Notes on *symmetric relations

- A relation can be neither symmetric or asymmetric
 - $R = \{ (a,b) \mid a=|b| \}$
 - This is not symmetric
 - -4 is not related to itself
 - This is not asymmetric
 - 4 is related to itself
 - Note that it is antisymmetric



Transitivity

- A relation is transitive if, for every $(a,b) \in R$ and $(b,c) \in R$, then $(a,c) \in R$
- If $a < b$ and $b < c$, then $a < c$
 - Thus, $<$ is transitive
- If $a = b$ and $b = c$, then $a = c$
 - Thus, $=$ is transitive



Transitivity examples

- Consider isAncestorOf()
 - Let Alice be Bob's parent, and Bob be Claire's parent
 - Thus, Alice is an ancestor of Bob, and Bob is an ancestor of Claire
 - Thus, Alice is an ancestor of Claire
 - Thus, isAncestorOf() is a transitive relation
- Consider isParentOf()
 - Let Alice be Bob's parent, and Bob be Claire's parent
 - Thus, Alice is a parent of Bob, and Bob is a parent of Claire
 - However, Alice is *not* a parent of Claire
 - Thus, isParentOf() is *not* a transitive relation



Relations of relations summary

	=	<	>	≤	≥
Reflexive	X			X	X
Irreflexive		X	X		
Symmetric	X				
Asymmetric		X	X		
Antisymmetric	X			X	X
Transitive	X	X	X	X	X



Combining relations

- There are two ways to combine relations R_1 and R_2
 - Via Boolean operators
 - Via relation “composition”



Combining relations via Boolean operators

- Consider two relations R_{\geq} and R_{\leq}
- We can combine them as follows:
 - $R_{\geq} \cup R_{\leq} = \text{all numbers } \geq \text{ OR } \leq$
 - That's all the numbers
 - $R_{\geq} \cap R_{\leq} = \text{all numbers } \geq \text{ AND } \leq$
 - That's all numbers equal to
 - $R_{\geq} \oplus R_{\leq} = \text{all numbers } \geq \text{ or } \leq, \text{ but not both}$
 - That's all numbers not equal to
 - $R_{\geq} - R_{\leq} = \text{all numbers } \geq \text{ that are not also } \leq$
 - That's all numbers strictly greater than
 - $R_{\leq} - R_{\geq} = \text{all numbers } \leq \text{ that are not also } \geq$
 - That's all numbers strictly less than
- Note that it's possible the result is the empty set



Combining relations via relational composition

- Let R be a relation from A to B , and S be a relation from B to C
 - Let $a \in A$, $b \in B$, and $c \in C$
 - Let $(a,b) \in R$, and $(b,c) \in S$
 - Then the composite of R and S consists of the ordered pairs (a,c)
 - We denote the relation by $S \circ R$
 - Note that S comes first when writing the composition!



Combining relations via relational composition

- Let M be the relation “is mother of”
- Let F be the relation “is father of”
- What is $M \circ F$?
 - If $(a,b) \in F$, then a is the father of b
 - If $(b,c) \in M$, then b is the mother of c
 - Thus, $M \circ F$ denotes the relation “maternal grandfather”
- What is $F \circ M$?
 - If $(a,b) \in M$, then a is the mother of b
 - If $(b,c) \in F$, then b is the father of c
 - Thus, $F \circ M$ denotes the relation “paternal grandmother”
- What is $M \circ M$?
 - If $(a,b) \in M$, then a is the mother of b
 - If $(b,c) \in M$, then b is the mother of c
 - Thus, $M \circ M$ denotes the relation “maternal grandmother”
- Note that M and F are not transitive relations!!!



Combining relations via relational composition

- Given relation R
 - $R \circ R$ can be denoted by R^2
 - $R^2 \circ R = (R \circ R) \circ R = R^3$
 - Example: M^3 is your mother's mother's mother



Representing Relations



In this slide set...

- Matrix review
- Two ways to represent relations
 - Via matrices
 - Via directed graphs



Matrix review

- We will only be dealing with zero-one matrices
 - Each element in the matrix is either a 0 or a 1

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 1 & 0 & 1 & 0 \end{bmatrix}$$

- These matrices will be used for Boolean operations
 - 1 is true, 0 is false



Matrix transposition

- Given a matrix \mathbf{M} , the transposition of \mathbf{M} , denoted \mathbf{M}^t , is the matrix obtained by switching the columns and rows of \mathbf{M}

$$\mathbf{M} = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix}$$

$$\mathbf{M}^t = \begin{bmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{bmatrix}$$

$$\mathbf{M} = \begin{bmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \\ 9 & 10 & 11 & 12 \\ 13 & 14 & 15 & 16 \end{bmatrix}$$

$$\mathbf{M}^t = \begin{bmatrix} 1 & 5 & 9 & 13 \\ 2 & 6 & 10 & 14 \\ 3 & 7 & 11 & 15 \\ 4 & 8 & 12 & 16 \end{bmatrix}$$

- In a “square” matrix, the main diagonal stays unchanged



Matrix join

- A *join* of two matrices performs a Boolean OR on each relative entry of the matrices
 - Matrices must be the same size
 - Denoted by the or symbol: \vee

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 1 & 0 & 1 & 0 \end{bmatrix} \vee \begin{bmatrix} 0 & 1 & 1 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 \\ 1 & 1 & 0 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 1 & 1 & 0 \\ 0 & 1 & 1 & 0 \\ 1 & 0 & 1 & 1 \\ 1 & 1 & 1 & 0 \end{bmatrix}$$



Matrix meet

- A *meet* of two matrices performs a Boolean AND on each relative entry of the matrices
 - Matrices must be the same size
 - Denoted by the or symbol: \wedge

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 1 & 0 & 1 & 0 \end{bmatrix} \wedge \begin{bmatrix} 0 & 1 & 1 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 \\ 1 & 1 & 0 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 \end{bmatrix}$$



Matrix Boolean product

- A *Boolean product* of two matrices is similar to matrix multiplication

$$c_{1,1} = a_{1,1} * b_{1,1} + a_{1,2} * b_{2,1} + a_{1,3} * b_{3,1} + a_{1,4} * b_{4,1}$$

- Instead of the sum of the products, it's the conjunction (and) of the disjunctions (ors)

$$c_{1,1} = a_{1,1} \wedge b_{1,1} \vee a_{1,2} \wedge b_{2,1} \vee a_{1,3} \wedge b_{3,1} \vee a_{1,4} \wedge b_{4,1}$$

- Denoted by the \odot symbol:

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 1 & 0 & 1 & 0 \end{bmatrix} \odot \begin{bmatrix} 0 & 1 & 1 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 \\ 1 & 1 & 0 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 1 & 1 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 1 & 1 & 1 \\ 0 & 1 & 1 & 1 \end{bmatrix}$$



Relations using matrices

- List the elements of sets A and B in a particular order
 - Order doesn't matter, but we'll generally use ascending order
- Create a matrix

$$\mathbf{M}_R = [m_{ij}]$$

$$m_{ij} = \begin{cases} 1 & \text{if } (a_i, b_j) \in R \\ 0 & \text{if } (a_i, b_j) \notin R \end{cases}$$



Relations using matrices

- Consider the relation of who is enrolled in which class
 - Let $A = \{ \text{Alice, Bob, Claire, Dan} \}$
 - Let $B = \{ \text{CS101, CS201, CS202} \}$
 - $R = \{ (a,b) \mid \text{person } a \text{ is enrolled in course } b \}$

	CS101	CS201	CS202
Alice	X		
Bob		X	X
Claire			
Dan		X	X

$$\mathbf{M}_R = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 0 \\ 0 & 1 & 1 \end{bmatrix}$$



Relations using matrices

- What is it good for?
 - It is how computers view relations
 - A 2-dimensional array
 - Very easy to view relationship properties
- We will generally consider relations on a single set
 - In other words, the domain and co-domain are the same set
 - And the matrix is square



Reflexivity

- Consider a reflexive relation: \leq
 - One which every element is related to itself
 - Let $A = \{ 1, 2, 3, 4, 5 \}$

$$\mathbf{M}_{\leq} = \begin{bmatrix} 1 & 1 & 1 & 1 & 1 \\ 0 & 1 & 1 & 1 & 1 \\ 0 & 0 & 1 & 1 & 1 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

If the center (main) diagonal is all 1's, a relation is reflexive



Irreflexivity

- Consider a reflexive relation: $<$
 - One which every element is *not* related to itself
 - Let $A = \{ 1, 2, 3, 4, 5 \}$

$$\mathbf{M}_{\leq} = \begin{bmatrix} 0 & 1 & 1 & 1 & 1 \\ 0 & 0 & 1 & 1 & 1 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

If the center (main) diagonal is all 0's, a relation is irreflexive



Symmetry

- Consider an symmetric relation R
 - One which if a is related to b then b is related to a for all (a,b)
 - Let $A = \{ 1, 2, 3, 4, 5 \}$

$$\mathbf{M}_{\leq} = \begin{bmatrix} 1 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 1 \end{bmatrix}$$

- If, for *every* value, it is the equal to the value in its transposed position, then the relation is symmetric



Asymmetry

- Consider an asymmetric relation: <
 - One which if a is related to b then b is *not* related to a for all (a,b)
 - Let $A = \{ 1, 2, 3, 4, 5 \}$

$$\mathbf{M}_{\leq} = \begin{bmatrix} 0 & 1 & 1 & 1 & 1 \\ 0 & 0 & 1 & 1 & 1 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

- If, for every value and the value in its transposed position, if they are not both 1, then the relation is asymmetric
- An asymmetric relation must also be irreflexive
- Thus, the main diagonal must be all 0's



Antisymmetry

- Consider an antisymmetric relation: \leq
 - One which if a is related to b then b is *not* related to a unless $a=b$ for all (a,b)
 - Let $A = \{ 1, 2, 3, 4, 5 \}$

$$\mathbf{M}_{\leq} = \begin{bmatrix} 1 & 1 & 1 & 1 & 1 \\ 0 & 1 & 1 & 1 & 1 \\ 0 & 0 & 1 & 1 & 1 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

- If, for every value and the value in its transposed position, if they are not both 1, then the relation is antisymmetric
- The center diagonal can have both 1's and 0's



Transitivity

- Consider an transitive relation: \leq
 - One which if a is related to b and b is related to c then a is related to c for all (a,b) , (b,c) and (a,c)
 - Let $A = \{ 1, 2, 3, 4, 5 \}$

$$\mathbf{M}_{\leq} = \begin{bmatrix} 1 & 1 & 1 & 1 & 1 \\ 0 & 1 & 1 & 1 & 1 \\ 0 & 0 & 1 & 1 & 1 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

- If, for every spot (a,b) and (b,c) that each have a 1, there is a 1 at (a,c) , then the relation is transitive
- Matrices don't show this property easily



Combining relations: via Boolean operators

- Let: $\mathbf{M}_R = \begin{bmatrix} 1 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}$ $\mathbf{M}_S = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 1 & 0 & 0 \end{bmatrix}$

- Join: $\mathbf{M}_{R \cup S} = \mathbf{M}_R \vee \mathbf{M}_S = \begin{bmatrix} 1 & 0 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 0 \end{bmatrix}$

- Meet: $\mathbf{M}_{R \cap S} = \mathbf{M}_R \wedge \mathbf{M}_S = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$



Combining relations: via relation composition

- Let:

$$\mathbf{M}_R = \begin{array}{c} a \\ b \\ c \end{array} \begin{array}{c} d \\ e \\ f \end{array} \begin{bmatrix} 1 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix} \quad \mathbf{M}_S = \begin{array}{c} d \\ e \\ f \end{array} \begin{array}{c} g \\ h \\ i \end{array} \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 1 & 0 & 0 \end{bmatrix}$$



- But why is this the case?

$$\mathbf{M}_{S \circ R} = \mathbf{M}_R \odot \mathbf{M}_S = \begin{array}{c} a \\ b \\ c \end{array} \begin{array}{c} g \\ h \\ i \end{array} \begin{bmatrix} 1 & 0 & 1 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$$



Representing relations using directed graphs


- A directed graph consists of:
 - A set V of vertices (or nodes)
 - A set E of edges (or arcs)
 - If (a, b) is in the relation, then there is an arrow from a to b
- Will generally use relations on a single set
- Consider our relation $R = \{ (a, b) \mid a \text{ divides } b \}$
- Old way:





Reflexivity

- Consider a reflexive relation: \leq
 - One which every element is related to itself
 - Let $A = \{ 1, 2, 3, 4, 5 \}$




If every node has a loop, a relation is reflexive



Irreflexivity

- Consider a reflexive relation: $<$
 - One which every element is *not* related to itself
 - Let $A = \{ 1, 2, 3, 4, 5 \}$




If every node does *not* have a loop, a relation is irreflexive



Symmetry

- Consider an symmetric relation R
 - One which if a is related to b then b is related to a for all (a,b)
 - Let $A = \{ 1, 2, 3, 4, 5 \}$




- If, for every edge, there is an edge in the other direction, then the relation is symmetric
- Loops are allowed, and do not need edges in the “other” direction

Note that this relation is neither reflexive nor irreflexive!

Asymmetry

- Consider an asymmetric relation: <
 - One which if a is related to b then b is *not* related to a for all (a,b)
 - Let $A = \{ 1, 2, 3, 4, 5 \}$




- A digraph is asymmetric if:
 1. If, for every edge, there is *not* an edge in the other direction, then the relation is asymmetric
 2. Loops are *not* allowed in an asymmetric digraph (recall it must be irreflexive)



Antisymmetry


- Consider an antisymmetric relation: \leq
 - One which if a is related to b then b is *not* related to a unless $a=b$ for all (a,b)
 - Let $A = \{ 1, 2, 3, 4, 5 \}$



- If, for every edge, there is *not* an edge in the other direction, then the relation is antisymmetric
- Loops are allowed in the digraph

Transitivity


- Consider an transitive relation: \leq
 - One which if a is related to b and b is related to c then a is related to c for all (a,b) , (b,c) and (a,c)
 - Let $A = \{ 1, 2, 3, 4, 5 \}$




- A digraph is transitive if, for there is a edge from a to c when there is a edge from a to b and from b to c

Sample questions

23.




24.




25.




26.



27.



28.



Which of the graphs are reflexive, irreflexive, symmetric, asymmetric, antisymmetric, or transitive

	23	24	25	26	27	28
Reflexive		Y		Y		Y
Irreflexive	Y		Y			
Symmetric					Y	Y
Asymmetric			Y			
Anti-symmetric		Y	Y			
Transitive						Y



Equivalence Relations



Introduction

- Certain combinations of relation properties are very useful
 - We won't have a chance to see many applications in this course
- In this set we will study equivalence relations
 - A relation that is reflexive, symmetric and transitive
- Next slide set we will study partial orderings
 - A relation that is reflexive, antisymmetric, and transitive
- The difference is whether the relation is symmetric or antisymmetric



Equivalence relations

- A relation on a set A is called an *equivalence relation* if it is reflexive, symmetric, and transitive
- Consider relation $R = \{ (a,b) \mid \text{len}(a) = \text{len}(b) \}$
 - Where $\text{len}(a)$ means the length of string a
 - It is reflexive: $\text{len}(a) = \text{len}(a)$
 - It is symmetric: if $\text{len}(a) = \text{len}(b)$, then $\text{len}(b) = \text{len}(a)$
 - It is transitive: if $\text{len}(a) = \text{len}(b)$ and $\text{len}(b) = \text{len}(c)$, then $\text{len}(a) = \text{len}(c)$
 - Thus, R is a equivalence relation



Equivalence relation example

- Consider the relation $R = \{ (a,b) \mid m \mid a-b \}$
 - Called “congruence modulo m ”
- Is it reflexive: $(a,a) \in R$ means that $m \mid a-a$
 - $a-a = 0$, which is divisible by m
- Is it symmetric: if $(a,b) \in R$ then $(b,a) \in R$
 - (a,b) means that $m \mid a-b$
 - Or that $km = a-b$. Negating that, we get $b-a = -km$
 - Thus, $m \mid b-a$, so $(b,a) \in R$
- Is it transitive: if $(a,b) \in R$ and $(b,c) \in R$ then $(a,c) \in R$
 - (a,b) means that $m \mid a-b$, or that $km = a-b$
 - (b,c) means that $m \mid b-c$, or that $lm = b-c$
 - (a,c) means that $m \mid a-c$, or that $nm = a-c$
 - Adding these two, we get $km+lm = (a-b) + (b-c)$
 - Or $(k+l)m = a-c$
 - Thus, m divides $a-c$, where $n = k+l$
- Thus, congruence modulo m is an equivalence relation



Sample questions

- Which of these relations on $\{0, 1, 2, 3\}$ are equivalence relations? Determine the properties of an equivalence relation that the others lack
- a) $\{ (0,0), (1,1), (2,2), (3,3) \}$
 - Has all the properties, thus, is an equivalence relation
- b) $\{ (0,0), (0,2), (2,0), (2,2), (2,3), (3,2), (3,3) \}$
 - Not reflexive: $(1,1)$ is missing
 - Not transitive: $(0,2)$ and $(2,3)$ are in the relation, but not $(0,3)$
- c) $\{ (0,0), (1,1), (1,2), (2,1), (2,2), (3,3) \}$
 - Has all the properties, thus, is an equivalence relation
- d) $\{ (0,0), (1,1), (1,3), (2,2), (2,3), (3,1), (3,2), (3,3) \}$
 - Not transitive: $(1,3)$ and $(3,2)$ are in the relation, but not $(1,2)$
- e) $\{ (0,0), (0,1), (0,2), (1,0), (1,1), (1,2), (2,0), (2,2), (3,3) \}$
 - Not symmetric: $(1,2)$ is present, but not $(2,1)$
 - Not transitive: $(2,0)$ and $(0,1)$ are in the relation, but not $(2,1)$