

# Programming and Numerical Analysis

30/01/2023

B. Haidar

Numerical Analysis



# Ordinary Differential Equations

# Learning Objectives of Lesson 1


---

- Recall basic definitions of ODEs:
  - Order
  - Linearity
  - Initial conditions
  - Solution
- Classify ODEs based on:
  - Order, linearity, and conditions.
- Classify the solution methods.

# Objectives

- Solve Ordinary Differential Equations (ODEs).
- Appreciate the importance of numerical methods in solving ODEs.
- Assess the reliability of the different techniques.
- Select the appropriate method for any particular problem.

# Derivatives



# Differential Equations

## Differential Equations

### Ordinary Differential Equations

$$\frac{d^2 v}{dt^2} + 6tv = 1$$

involve one or more  
Ordinary derivatives of  
unknown functions

### Partial Differential Equations

$$\frac{\partial^2 u}{\partial y^2} - \frac{\partial^2 u}{\partial x^2} = 0$$

involve one or more  
partial derivatives of  
unknown functions

# Ordinary Differential Equations

**Ordinary Differential Equations (ODEs)** involve one or more ordinary derivatives of unknown functions with respect to one independent variable

*Examples :*

$$\frac{dv(t)}{dt} - v(t) = e^t$$

x(t): unknown function

$$\frac{d^2 x(t)}{dt^2} - 5 \frac{dx(t)}{dt} + 2x(t) = \cos(t)$$

t: independent variable

# Example of ODE - Model of Falling Parachutist


The velocity of a falling parachutist is given by:

$$\frac{dv}{dt} = 9.8 - \frac{c}{M}v$$

$M$  : mass

$c$  : drag coefficient

$v$  : velocity





# Definitions

$$\frac{dv}{dt} = 9.8 - \frac{c}{M} v$$

Ordinary  
differential  
equation

# Definitions (Cont.)

$$\frac{dv}{dt} = 9.8 - \frac{c}{M}v$$

(Dependent variable)  
unknown  
function to be  
determined

# Definitions (Cont.)

$$\frac{dv}{dt} = 9.8 - \frac{c}{M} v$$

(independent variable)  
the variable with respect to which  
other variables are differentiated

# Order of Differential Equation

The **order** of an ordinary differential equations is the order of the highest order derivative.

*Examples :*

$$\frac{dx(t)}{dt} - x(t) = e^t$$

First order ODE

$$\frac{d^2 x(t)}{dt^2} - 5 \frac{dx(t)}{dt} + 2x(t) = \cos(t)$$

Second order ODE

$$\left( \frac{d^2 x(t)}{dt^2} \right)^3 - \frac{dx(t)}{dt} + 2x^4(t) = 1$$

Second order ODE

# Solution of Differential Equation

A **solution** to a differential equation is a function that satisfies the equation.

*Example :*

$$\frac{dx(t)}{dt} + x(t) = 0$$

*Solution*  $x(t) = e^{-t}$

Proof :

$$\frac{dx(t)}{dt} = -e^{-t}$$

$$\frac{dx(t)}{dt} + x(t) = -e^{-t} + e^{-t} = 0$$

# Linear ODE

An ODE is linear if

The unknown function and its derivatives appear to power one

No product of the unknown function and/or its derivatives

*Examples :*

$$\frac{dx(t)}{dt} - x(t) = e^t$$

Linear ODE

$$\frac{d^2x(t)}{dt^2} - 5\frac{dx(t)}{dt} + 2t^2x(t) = \cos(t)$$

Linear ODE

$$\left(\frac{d^2x(t)}{dt^2}\right)^3 - \frac{dx(t)}{dt} + \sqrt{x(t)} = 1$$

Non-linear ODE

# Nonlinear ODE

An ODE is linear if

The unknown function and its derivatives appear to power one

No product of the unknown function and/or its derivatives

Examples of nonlinear ODE :

$$\frac{dx(t)}{dt} - \cos(x(t)) = 1$$

$$\frac{d^2x(t)}{dt^2} - 5 \frac{dx(t)}{dt} x(t) = 2$$

$$\frac{d^2x(t)}{dt^2} - \left| \frac{dx(t)}{dt} \right| + x(t) = 1$$

# Solution of Ordinary Differential Equations

$$x(t) = \cos(2t)$$

is a solution to the ODE

$$\frac{d^2 x(t)}{dt^2} + 4x(t) = 0$$

Is it unique?

All functions of the form  $x(t) = \cos(2t + c)$   
(where  $c$  is a real constant) are solutions.



# Uniqueness of a Solution

In order to uniquely specify a solution to an  $n^{\text{th}}$  order differential equation we need  $n$  conditions.

$$\frac{d^2 y(x)}{dx^2} + 4y(x) = 0$$

Second order ODE

$$y(0) = a$$

$$\dot{y}(0) = b$$

Two conditions are needed to uniquely specify the solution

# Auxiliary Conditions

## Auxiliary Conditions

### Initial Conditions

- All conditions are at **one point of the independent variable**

### Boundary Conditions

- The conditions are **not at one point of the independent variable**

# Boundary Value and Initial Value Problems

30/01/2023

B. Haidar

Numerical Analysis

## Initial-Value Problems

- The auxiliary conditions are at **one point of the independent variable**

$$\ddot{x} + 2\dot{x} + x = e^{-2t}$$

$$x(0) = 1, \dot{x}(0) = 2.5$$

same

## Boundary-Value Problems

- The auxiliary conditions are **not at one point of the independent variable**
- More difficult to solve than initial value problems

$$\ddot{x} + 2\dot{x} + x = e^{-2t}$$

$$x(0) = 1, x(2) = 1.5$$

different

# Classification of ODEs

ODEs can be classified in different ways:

- Order
  - First order ODE
  - Second order ODE
  - $N^{\text{th}}$  order ODE
- Linearity
  - Linear ODE
  - Nonlinear ODE
- Auxiliary conditions
  - Initial value problems
  - Boundary value problems


# Analytical Solutions

- Analytical Solutions to ODEs are available for linear ODEs and special classes of nonlinear differential equations.

# Numerical Solutions

- Numerical methods are used to obtain a graph or a table of the unknown function.
- Most of the Numerical methods used to solve ODEs are based directly (or indirectly) on the truncated Taylor series expansion.

# Classification of the Methods



# Taylor Series Methods



# Learning Objectives of Lesson 2

---

- ❑ Derive Euler formula using the Taylor series expansion.
- ❑ Solve the first order ODEs using Euler method.
- ❑ Assess the error level when using Euler method.
- ❑ Appreciate different types of errors in the numerical solution of ODEs.
- ❑ Improve Euler method using higher-order Taylor Series.

# Taylor Series Method

The problem to be solved is a first order ODE:

$$\frac{dy(x)}{dx} = f(x, y), \quad y(x_0) = y_0$$

Estimates of the solution at different base points:

$$y(x_0 + h), \quad y(x_0 + 2h), \quad y(x_0 + 3h), \quad \dots$$

are computed using the truncated Taylor series expansions.

# Taylor Series Expansion

## Truncated Taylor Series Expansion

$$y(x_0 + h) \approx \sum_{k=0}^n \frac{h^k}{k!} \left( \frac{d^k y}{dx^k} \Big|_{x=x_0, y=y_0} \right)$$
$$\approx y(x_0) + h \frac{dy}{dx} \Big|_{x=x_0, y=y_0} + \frac{h^2}{2!} \frac{d^2 y}{dx^2} \Big|_{x=x_0, y=y_0} + \dots + \frac{h^n}{n!} \frac{d^n y}{dx^n} \Big|_{x=x_0, y=y_0}$$

The  $n^{\text{th}}$  order Taylor series method uses the  $n^{\text{th}}$  order Truncated Taylor series expansion.

# Euler Method

- First order Taylor series method is known as Euler Method.
- Only the constant term and linear term are used in the Euler method.
- The error due to the use of the truncated Taylor series is of order  $O(h^2)$ .

# First Order Taylor Series Method (Euler Method)

$$y(x_0 + h) = y(x_0) + h \left. \frac{dy}{dx} \right|_{\substack{x=x_0, \\ y=y_0}} + O(h^2)$$

*Notation :*

$$x_n = x_0 + nh, \quad y_n = y(x_n),$$

$$\left. \frac{dy}{dx} \right|_{\substack{x=x_i, \\ y=y_i}} = f(x_i, y_i)$$

*Euler Method*

$$y_{i+1} = y_i + h f(x_i, y_i)$$

# Euler Method

Problem :

Given the first order ODE :  $\dot{y}(x) = f(x, y)$

with the initial condition :  $y_0 = y(x_0)$


Determine :  $y_i = y(x_0 + ih)$  for  $i = 1, 2, \dots$

Euler Method :


$$y_0 = y(x_0)$$

$$y_{i+1} = y_i + h f(x_i, y_i) \quad \text{for } i = 1, 2, \dots$$

# Interpretation of Euler Method



# Interpretation of Euler Method





# Example 1

Use Euler method to solve the ODE:

$$\frac{dy}{dx} = 1 + x^2, \quad y(1) = -4$$

to determine  $y(1.01)$ ,  $y(1.02)$  and  $y(1.03)$ .

# Example 1

$$f(x, y) = 1 + x^2, \quad x_0 = 1, \quad y_0 = -4, \quad h = 0.01$$

Euler Method

$$y_{i+1} = y_i + h f(x_i, y_i)$$

$$\text{Step1: } y_1 = y_0 + h f(x_0, y_0) = -4 + 0.01(1 + (1)^2) = -3.98$$

$$\text{Step2: } y_2 = y_1 + h f(x_1, y_1) = -3.98 + 0.01(1 + (1.01)^2) = -3.9598$$

$$\text{Step3: } y_3 = y_2 + h f(x_2, y_2) = -3.9598 + 0.01(1 + (1.02)^2) = -3.9394$$

# Example 1

$$f(x, y) = 1 + x^2, \quad x_0 = 1, \quad y_0 = -4, \quad h = 0.01$$

Summary of the result:

i	$x_i$	$y_i$
0	1.00	-4.00
1	1.01	-3.98
2	1.02	-3.9595
3	1.03	-3.9394

# Example 1

$$f(x, y) = 1 + x^2, \quad x_0 = 1, \quad y_0 = -4, \quad h = 0.01$$

Comparison with true value:

i	$x_i$	$y_i$	True value of $y_i$
0	1.00	-4.00	-4.00
1	1.01	-3.98	-3.97990
2	1.02	-3.9595	-3.95959
3	1.03	-3.9394	-3.93909