



Calculus 1

Dr. Yamar Hamwi

Al-Manara University

2024-2025

<https://manara.edu.sy/>

Calculus 1

Lecture 10

Integrals

Chapter 5

Integrals

5.7 The Definite integral

5.8 The Fundamental Theorem of Calculus

5.9 Definite integral Substitutions

5.10 Evaluating Definite Integrals by Parts


5.11 Areas Between Curves

5.12 Solids of Revolution: the Disk Method

5.13 Volume by Washers for Rotation About the x-Axis

5.14 Length of a Curve $y = f(x)$

The Definite integral



The diagram illustrates the components of a definite integral $\int_a^b f(x) dx$. The integral sign $\int$ is labeled as the "Integral sign". The upper limit b is labeled as the "Upper limit of integration". The lower limit a is labeled as the "Lower limit of integration". The function $f(x)$ is labeled as "The function $f(x)$ is the integrand.". The variable x in dx is labeled as " x is the variable of integration.". A bracket under the entire expression $\int_a^b f(x) dx$ is labeled as "Integral of f from a to b ". A separate label points to this bracketed expression, stating "When you find the value of the integral, you have evaluated the integral."

Upper limit of integration

Integral sign

Lower limit of integration

The function $f(x)$ is the integrand.

x is the variable of integration.


Integral of f from a to b

When you find the value of the integral, you have evaluated the integral.

THEOREM 4.5 The Definite Integral as the Area of a Region

If f is continuous and nonnegative on the closed interval $[a, b]$, then the area of the region bounded by the graph of f , the x -axis, and the vertical lines $x = a$ and $x = b$ is

$$\text{Area} = \int_a^b f(x) \, dx.$$



rules satisfied by definite integrals

1. *Order of Integration:* $\int_b^a f(x) dx = -\int_a^b f(x) dx$

A definition

2. *Zero Width Interval:* $\int_a^a f(x) dx = 0$

A definition
when $f(a)$ exists

3. *Constant Multiple:* $\int_a^b kf(x) dx = k \int_a^b f(x) dx$

Any constant k

4. *Sum and Difference:* $\int_a^b (f(x) \pm g(x)) dx = \int_a^b f(x) dx \pm \int_a^b g(x) dx$

5. *Additivity:* $\int_a^b f(x) dx + \int_b^c f(x) dx = \int_a^c f(x) dx$

6. *Domination:* If $f(x) \geq g(x)$ on $[a, b]$ then $\int_a^b f(x) dx \geq \int_a^b g(x) dx$.

If $f(x) \geq 0$ on $[a, b]$ then $\int_a^b f(x) dx \geq 0$.

Special case

The Fundamental Theorem of Calculus

THEOREM 7.1 Continuity Implies Integrability

If a function f is continuous on the closed interval $[a, b]$, then f is integrable on $[a, b]$. That is, $\int_a^b f(x) dx$ exists.

THEOREM 7.2 The Fundamental Theorem of Calculus

If a function f is continuous on the closed interval $[a, b]$ and F is an antiderivative of f on the interval $[a, b]$, then

$$\int_a^b f(x) dx = F(b) - F(a).$$

1. Find an antiderivative F of f , and
2. Calculate the number $F(b) - F(a)$, which is equal to $\int_a^b f(x) dx$.

The Fundamental Theorem of Calculus

$$\begin{aligned} \text{(a)} \quad \int_0^{\pi} \cos x \, dx &= \sin x \Big|_0^{\pi} \\ &= \sin \pi - \sin 0 = 0 - 0 = 0 \end{aligned}$$

$$\begin{aligned} \text{(b)} \quad \int_{-\pi/4}^0 \sec x \tan x \, dx &= \sec x \Big|_{-\pi/4}^0 \\ &= \sec 0 - \sec \left(-\frac{\pi}{4} \right) = 1 - \sqrt{2} \end{aligned}$$

$$\frac{d}{dx} \sec x = \sec x \tan x$$

$$\begin{aligned} \text{(c)} \quad \int_1^4 \left(\frac{3}{2} \sqrt{x} - \frac{4}{x^2} \right) dx &= \left[x^{3/2} + \frac{4}{x} \right]_1^4 \\ &= \left[(4)^{3/2} + \frac{4}{4} \right] - \left[(1)^{3/2} + \frac{4}{1} \right] \\ &= [8 + 1] - [5] = 4 \end{aligned}$$

$$\frac{d}{dx} \left(x^{3/2} + \frac{4}{x} \right) = \frac{3}{2} x^{1/2} - \frac{4}{x^2}$$



The Fundamental Theorem of Calculus

EXAMPLE


Evaluate $\int_0^2 |2x - 1| dx$.

Solution

$$|2x - 1| = \begin{cases} -(2x - 1), & x < \frac{1}{2} \\ 2x - 1, & x \geq \frac{1}{2} \end{cases}$$

From this, you can rewrite the integral in two parts.

$$\begin{aligned} \int_0^2 |2x - 1| dx &= \int_0^{1/2} -(2x - 1) dx + \int_{1/2}^2 (2x - 1) dx \\ &= \left[-x^2 + x \right]_0^{1/2} + \left[x^2 - x \right]_{1/2}^2 \\ &= \left(-\frac{1}{4} + \frac{1}{2} \right) - (0 + 0) + (4 - 2) - \left(\frac{1}{4} - \frac{1}{2} \right) \\ &= \frac{5}{2} \end{aligned}$$



The definite integral of y on $[0, 2]$ is $\frac{5}{2}$.

Using the Fundamental Theorem to Find Area

Find the area of the region bounded by the graph of

$$y = 2x^2 - 3x + 2$$

the x -axis, and the vertical lines $x = 0$ and $x = 2$, as shown in Figure 4.29.

Solution Note that $y > 0$ on the interval $[0, 2]$.


$$\begin{aligned}\text{Area} &= \int_0^2 (2x^2 - 3x + 2) \, dx \\ &= \left[\frac{2x^3}{3} - \frac{3x^2}{2} + 2x \right]_0^2 \\ &= \left(\frac{16}{3} - 6 + 4 \right) - (0 - 0 + 0) \\ &= \frac{10}{3}\end{aligned}$$

Integrate between $x = 0$ and $x = 2$.

Find antiderivative.

Apply Fundamental Theorem.

Simplify.



THEOREM 7 – Substitution in Definite Integrals

If g' is continuous on the interval $[a, b]$ and f is continuous on the range of $g(x) = u$, then

$$\int_a^b f(g(x)) \cdot g'(x) dx = \int_{g(a)}^{g(b)} f(u) du.$$

EXAMPLE 1 Evaluate $\int_{-1}^1 3x^2 \sqrt{x^3 + 1} dx$.

Definite integral Substitutions

$$\int_{-1}^1 3x^2 \sqrt{x^3 + 1} dx$$

$$= \int_0^2 \sqrt{u} du$$

$$= \left. \frac{2}{3} u^{3/2} \right|_0^2$$

$$= \frac{2}{3} [2^{3/2} - 0^{3/2}] = \frac{2}{3} [2\sqrt{2}] = \frac{4\sqrt{2}}{3}$$

Let $u = x^3 + 1$, $du = 3x^2 dx$.

When $x = -1$, $u = (-1)^3 + 1 = 0$.

When $x = 1$, $u = (1)^3 + 1 = 2$.

Evaluate the new definite integral.



Definite integral Substitutions

$$\int_{-\pi/4}^{\pi/4} \tan x \, dx$$

$$\int_{-\pi/4}^{\pi/4} \tan x \, dx = \int_{-\pi/4}^{\pi/4} \frac{\sin x}{\cos x} \, dx$$

$$= - \int_{\sqrt{2}/2}^{\sqrt{2}/2} \frac{du}{u}$$

$$= -\ln |u| \Big|_{\sqrt{2}/2}^{\sqrt{2}/2} = 0$$

Let $u = \cos x$, $du = -\sin x \, dx$.

When $x = -\pi/4$, $u = \sqrt{2}/2$.

When $x = \pi/4$, $u = \sqrt{2}/2$.

Integrate, zero width interval

Definite integral Substitutions

$$\int_0^{\pi/2} \frac{2 \sin x \cos x}{(1 + \sin^2 x)^3} dx$$

$$\begin{aligned} \int_0^{\pi/2} \frac{2 \sin x \cos x}{(1 + \sin^2 x)^3} dx &= \int_1^2 \frac{1}{u^3} du \\ &= -\frac{1}{2u^2} \Bigg|_1^2 \\ &= -\frac{1}{8} - \left(-\frac{1}{2}\right) = \frac{3}{8} \end{aligned}$$

Let $u = 1 + \sin^2 x$, $du = 2 \sin x \cos x dx$.
When $x = 0$, $u = 1$.
When $x = \pi/2$, $u = 2$.



Definite integral Substitutions

$$\int_3^5 \frac{2x - 3}{\sqrt{x^2 - 3x + 1}} dx.$$

$$\int_3^5 \frac{2x - 3}{\sqrt{x^2 - 3x + 1}} dx = \int_1^{11} \frac{du}{\sqrt{u}}$$

$$u = x^2 - 3x + 1, du = (2x - 3) dx;$$

$$u = 1 \text{ when } x = 3, u = 11 \text{ when } x = 5$$

$$= \int_1^{11} u^{-1/2} du$$

$$= 2\sqrt{u} \Big|_1^{11} = 2(\sqrt{11} - 1) \approx 4.63. \quad \text{Table 8.1, Formula 2}$$





Evaluating Definite Integrals by Parts

Integration by Parts Formula for Definite Integrals

$$\int_a^b f(x)g'(x) dx = f(x)g(x) \Big|_a^b - \int_a^b f'(x)g(x) dx$$

Evaluating Definite Integrals by Parts


EXAMPLE 6 Find the area of the region bounded by the curve $y = xe^{-x}$ and the x -axis from $x = 0$ to $x = 4$.

Solution The region is shaded in Figure 8.1. Its area is

$$\int_0^4 xe^{-x} dx.$$

Let $u = x$, $dv = e^{-x} dx$, $v = -e^{-x}$, and $du = dx$. Then,

$$\begin{aligned} \int_0^4 xe^{-x} dx &= -xe^{-x} \Big|_0^4 - \int_0^4 (-e^{-x}) dx \\ &= [-4e^{-4} - (-0e^{-0})] + \int_0^4 e^{-x} dx \\ &= -4e^{-4} - e^{-x} \Big|_0^4 \\ &= -4e^{-4} - (e^{-4} - e^{-0}) = 1 - 5e^{-4} \approx \end{aligned}$$



The Definite integral

$$\int_0^{\pi/4} \sqrt{1 + \cos 4x} \, dx.$$

Solution To eliminate the square root, we use the identity

$$\cos^2 \theta = \frac{1 + \cos 2\theta}{2} \quad \text{or} \quad 1 + \cos 2\theta = 2 \cos^2 \theta.$$

With $\theta = 2x$, this becomes

$$1 + \cos 4x = 2 \cos^2 2x.$$

Therefore,

$$\begin{aligned} \int_0^{\pi/4} \sqrt{1 + \cos 4x} \, dx &= \int_0^{\pi/4} \sqrt{2 \cos^2 2x} \, dx = \int_0^{\pi/4} \sqrt{2} \sqrt{\cos^2 2x} \, dx \\ &= \sqrt{2} \int_0^{\pi/4} |\cos 2x| \, dx = \sqrt{2} \int_0^{\pi/4} \cos 2x \, dx \\ &= \sqrt{2} \left[\frac{\sin 2x}{2} \right]_0^{\pi/4} = \frac{\sqrt{2}}{2} [1 - 0] = \frac{\sqrt{2}}{2}. \end{aligned}$$

$\cos 2x \geq 0$ on
 $[0, \pi/4]$



The Definite integral

$$\int_0^{\pi/2} \sin^2 2\theta \cos^3 2\theta d\theta$$

$$\begin{aligned}\int_0^{\pi/2} \sin^2 2\theta \cos^3 2\theta d\theta &= \int_0^{\pi/2} \sin^2 2\theta (1 - \sin^2 2\theta) \cos 2\theta d\theta \\ &= \int_0^{\pi/2} \sin^2 2\theta \cos 2\theta d\theta - \int_0^{\pi/2} \sin^4 2\theta \cos 2\theta d\theta = \left[\frac{1}{2} \cdot \frac{\sin^3 2\theta}{3} - \frac{1}{2} \cdot \frac{\sin^5 2\theta}{5} \right]_0^{\pi/2} = 0\end{aligned}$$

The Definite integral

Evaluate $\int_{\pi/6}^{\pi/3} \frac{\cos^3 x}{\sqrt{\sin x}} dx.$

$$\begin{aligned}\int_{\pi/6}^{\pi/3} \frac{\cos^3 x}{\sqrt{\sin x}} dx &= \int_{\pi/6}^{\pi/3} \frac{\cos^2 x \cos x}{\sqrt{\sin x}} dx \\&= \int_{\pi/6}^{\pi/3} \frac{(1 - \sin^2 x)(\cos x)}{\sqrt{\sin x}} dx \\&= \int_{\pi/6}^{\pi/3} [(\sin x)^{-1/2} - (\sin x)^{3/2}] \cos x dx \\&= \left[\frac{(\sin x)^{1/2}}{1/2} - \frac{(\sin x)^{5/2}}{5/2} \right]_{\pi/6}^{\pi/3} \\&= 2 \left(\frac{\sqrt{3}}{2} \right)^{1/2} - \frac{2}{5} \left(\frac{\sqrt{3}}{2} \right)^{5/2} - \sqrt{2} + \frac{\sqrt{32}}{80}\end{aligned}$$

The Definite integral

Evaluate $\int_0^2 \frac{x}{1 + e^{-x^2}} dx$.

$$\int \frac{du}{1 + e^u} = u - \ln(1 + e^u) + C.$$

Example 1.21

Let $u = -x^2$. Then $du = -2x dx$, and you have

$$\begin{aligned} \int \frac{x}{1 + e^{-x^2}} dx &= -\frac{1}{2} \int \frac{-2x dx}{1 + e^{-x^2}} \\ &= -\frac{1}{2} [-x^2 - \ln(1 + e^{-x^2})] + C \\ &= \frac{1}{2} [x^2 + \ln(1 + e^{-x^2})] + C. \end{aligned}$$

So, the value of the definite integral is

$$\int_0^2 \frac{x}{1 + e^{-x^2}} dx = \frac{1}{2} \left[x^2 + \ln(1 + e^{-x^2}) \right]_0^2 = \frac{1}{2} [4 + \ln(1 + e^{-4}) - \ln 2]$$



Definite Integrals of Symmetric Functions




FIGURE : (a) For f an even function, the integral from $-a$ to a is twice the integral from 0 to a . (b) For f an odd function, the integral from $-a$ to a equals 0.

Definite Integrals of Symmetric Functions

THEOREM Let f be continuous on the symmetric interval $[-a, a]$.

(a) If f is even, then $\int_{-a}^a f(x) dx = 2 \int_0^a f(x) dx$.

(b) If f is odd, then $\int_{-a}^a f(x) dx = 0$.

EXAMPLE

Evaluate $\int_{-2}^2 (x^4 - 4x^2 + 6) dx$.

Definite Integrals of Symmetric Functions

Solution Since $f(x) = x^4 - 4x^2 + 6$ satisfies $f(-x) = f(x)$, it is even on the symmetric interval $[-2, 2]$, so


$$\begin{aligned}\int_{-2}^2 (x^4 - 4x^2 + 6) dx &= 2 \int_0^2 (x^4 - 4x^2 + 6) dx \\ &= 2 \left[\frac{x^5}{5} - \frac{4}{3}x^3 + 6x \right]_0^2 \\ &= 2 \left(\frac{32}{5} - \frac{32}{3} + 12 \right) = \frac{232}{15}.\end{aligned}$$



Areas Between Curves

DEFINITION If f and g are continuous with $f(x) \geq g(x)$ throughout $[a, b]$, then the **area of the region between the curves $y = f(x)$ and $y = g(x)$ from a to b** is the integral of $(f - g)$ from a to b :

$$A = \int_a^b [f(x) - g(x)] dx.$$




Areas Between Curves

EXAMPLE 4 Find the area of the region enclosed by the parabola $y = 2 - x^2$ and the line $y = -x$.

Solution First we sketch the two curves (Figure 5.28). The limits of integration are found by solving $y = 2 - x^2$ and $y = -x$ simultaneously for x .

$$\begin{aligned}2 - x^2 &= -x && \text{Equate } f(x) \text{ and } g(x). \\x^2 - x - 2 &= 0 && \text{Rewrite.} \\(x + 1)(x - 2) &= 0 && \text{Factor.} \\x = -1, \quad x = 2. &&& \text{Solve.}\end{aligned}$$

The region runs from $x = -1$ to $x = 2$. The limits of integration are $a = -1$, $b = 2$.



Areas Between Curves

The area between the curves is

$$\begin{aligned} A &= \int_a^b [f(x) - g(x)] dx = \int_{-1}^2 [(2 - x^2) - (-x)] dx \\ &= \int_{-1}^2 (2 + x - x^2) dx = \left[2x + \frac{x^2}{2} - \frac{x^3}{3} \right]_{-1}^2 \\ &= \left(4 + \frac{4}{2} - \frac{8}{3} \right) - \left(-2 + \frac{1}{2} + \frac{1}{3} \right) = \frac{9}{2}. \end{aligned}$$

Areas Between Curves

EXAMPLE 5 Find the area of the region in the first quadrant that is bounded above by $y = \sqrt{x}$ and below by the x -axis and the line $y = x - 2$.

Solution

$$\sqrt{x} = x - 2$$

$$x = (x - 2)^2 = x^2 - 4x + 4$$

$$x^2 - 5x + 4 = 0$$

$$(x - 1)(x - 4) = 0$$

$$x = 1, \quad x = 4.$$


Equate $f(x)$ and $g(x)$.

Square both sides.

Rewrite.

Factor.

Solve.



Areas Between Curves

$$\text{For } 0 \leq x \leq 2: \quad f(x) - g(x) = \sqrt{x} - 0 = \sqrt{x}$$

$$\text{For } 2 \leq x \leq 4: \quad f(x) - g(x) = \sqrt{x} - (x - 2) = \sqrt{x} - x + 2$$

$$\text{Total area} = \underbrace{\int_0^2 \sqrt{x} \, dx}_{\text{area of A}} + \underbrace{\int_2^4 (\sqrt{x} - x + 2) \, dx}_{\text{area of B}}$$

$$= \left[\frac{2}{3} x^{3/2} \right]_0^2 + \left[\frac{2}{3} x^{3/2} - \frac{x^2}{2} + 2x \right]_2^4$$


$$= \frac{2}{3} (2)^{3/2} - 0 + \left(\frac{2}{3} (4)^{3/2} - 8 + 8 \right) - \left(\frac{2}{3} (2)^{3/2} - 2 + 4 \right)$$

$$= \frac{2}{3} (8) - 2 = \frac{10}{3}.$$

Volumes

DEFINITION The **volume** of a solid of integrable cross-sectional area $A(x)$ from $x = a$ to $x = b$ is the integral of A from a to b ,


$$V = \int_a^b A(x) dx.$$



Solids of Revolution: the Disk Method

Volume by Disks for Rotation About the x -Axis

$$V = \int_a^b A(x) dx = \int_a^b \pi [R(x)]^2 dx.$$



Horizontal axis of revolution



Solids of Revolution: the Disk Method

EXAMPLE

Find the volume of the solid formed by revolving the region bounded by the graph of

$$f(x) = \sqrt{\sin x}$$

and the x -axis ($0 \leq x \leq \pi$) about the x -axis.





Solids of Revolution: the Disk Method

Solution From the representative rectangle in the upper graph in Figure , you can see that the radius of this solid is

$$\begin{aligned} R(x) &= f(x) \\ &= \sqrt{\sin x}. \end{aligned}$$

So, the volume of the solid of revolution is

$$V = \pi \int_a^b [R(x)]^2 dx = \pi \int_0^{\pi} (\sqrt{\sin x})^2 dx$$

Apply disk method.

$$= \pi \int_0^{\pi} \sin x dx$$

Simplify.

$$= \pi \left[-\cos x \right]_0^{\pi}$$

Integrate.

$$= \pi(1 + 1)$$

$$= 2\pi.$$



Solids of Revolution: the Disk Method

EXAMPLE

Find the volume of the solid formed by revolving the region bounded by

$$f(x) = 2 - x^2$$

and $g(x) = 1$ about the line $y = 1$, as shown in Figure





Solids of Revolution: the Disk Method

Solution By equating $f(x)$ and $g(x)$, you can determine that the two graphs intersect when $x = \pm 1$. To find the radius, subtract $g(x)$ from $f(x)$.

$$\begin{aligned} R(x) &= f(x) - g(x) \\ &= (2 - x^2) - 1 \\ &= 1 - x^2 \end{aligned}$$


Finally, integrate between -1 and 1 to find the volume.

$$\begin{aligned} V &= \pi \int_a^b [R(x)]^2 dx = \pi \int_{-1}^1 (1 - x^2)^2 dx && \text{Apply disk method.} \\ &= \pi \int_{-1}^1 (1 - 2x^2 + x^4) dx && \text{Simplify.} \\ &= \pi \left[x - \frac{2x^3}{3} + \frac{x^5}{5} \right]_{-1}^1 && \text{Integrate.} \\ &= \frac{16\pi}{15} \end{aligned}$$

Volume by Washers for Rotation About the x-Axis

Volume by Washers for Rotation About the x-Axis



$$V = \int_a^b A(x) dx = \int_a^b \pi ([R(x)]^2 - [r(x)]^2) dx.$$



Volume by Washers for Rotation About the x-Axis

EXAMPLE

Find the volume of the solid formed by revolving the region bounded by the graphs of $y = \sqrt{x}$ and $y = x^2$ about the x -axis, as shown in Figure



Volume by Washers for Rotation About the x-Axis

Solution In Figure , you can see that the outer and inner radii are as follows.

$$R(x) = \sqrt{x}$$

Outer radius

$$r(x) = x^2$$

Inner radius

Integrating between 0 and 1 produces

$$V = \pi \int_a^b ([R(x)]^2 - [r(x)]^2) dx$$

Apply washer method.

$$= \pi \int_0^1 [(\sqrt{x})^2 - (x^2)^2] dx$$

$$= \pi \int_0^1 (x - x^4) dx$$

Simplify.

$$= \pi \left[\frac{x^2}{2} - \frac{x^5}{5} \right]_0^1$$


Integrate.

$$= \frac{3\pi}{10}.$$

Length of a Curve $y = f(x)$

DEFINITION If f' is continuous on $[a, b]$, then the **length (arc length)** of the curve $y = f(x)$ from the point $A = (a, f(a))$ to the point $B = (b, f(b))$ is the value of the integral

$$L = \int_a^b \sqrt{1 + [f'(x)]^2} dx = \int_a^b \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx. \quad (3)$$



Length of a Curve $y = f(x)$

Find the length of the curve

$$y = \frac{4\sqrt{2}}{3}x^{3/2} - 1, \quad 0 \leq x \leq 1.$$


Solution We use Equation (3) with $a = 0$, $b = 1$, and

$$y = \frac{4\sqrt{2}}{3}x^{3/2} - 1$$

$$\frac{dy}{dx} = \frac{4\sqrt{2}}{3} \cdot \frac{3}{2}x^{1/2} = 2\sqrt{2}x^{1/2}$$

$$\left(\frac{dy}{dx}\right)^2 = (2\sqrt{2}x^{1/2})^2 = 8x.$$

$$x = 1, y \approx 0.89$$



Length of a Curve $y = f(x)$

The length of the curve over $x = 0$ to $x = 1$ is

$$\begin{aligned} L &= \int_0^1 \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx = \int_0^1 \sqrt{1 + 8x} dx \\ &= \frac{2}{3} \cdot \frac{1}{8} (1 + 8x)^{3/2} \Big|_0^1 = \frac{13}{6} \approx 2.17. \end{aligned}$$

Eq. (3) with
 $a = 0, b = 1$.

Let $u = 1 + 8x$,
integrate, and
replace u by
 $1 + 8x$.

EXAMPLE 2 Find the length of the graph of

$$f(x) = \frac{x^3}{12} + \frac{1}{x}, \quad 1 \leq x \leq 4.$$

Length of a Curve $y = f(x)$

Solution A graph of the function is shown in Figure . To use Equation (3), we find


$$f'(x) = \frac{x^2}{4} - \frac{1}{x^2}$$

so

$$\begin{aligned} 1 + [f'(x)]^2 &= 1 + \left(\frac{x^2}{4} - \frac{1}{x^2}\right)^2 = 1 + \left(\frac{x^4}{16} - \frac{1}{2} + \frac{1}{x^4}\right) \\ &= \frac{x^4}{16} + \frac{1}{2} + \frac{1}{x^4} = \left(\frac{x^2}{4} + \frac{1}{x^2}\right)^2. \end{aligned}$$

The length of the graph over $[1, 4]$ is

$$\begin{aligned} L &= \int_1^4 \sqrt{1 + [f'(x)]^2} dx = \int_1^4 \left(\frac{x^2}{4} + \frac{1}{x^2}\right) dx \\ &= \left[\frac{x^3}{12} - \frac{1}{x}\right]_1^4 = \left(\frac{64}{12} - \frac{1}{4}\right) - \left(\frac{1}{12} - 1\right) = \frac{72}{12} = 6. \end{aligned}$$



Length of a Curve $y = f(x)$

EXAMPLE Find the length of the curve

$$y = \frac{1}{2} (e^x + e^{-x}), \quad 0 \leq x \leq 2.$$

Solution We use Equation (3) with $a = 0$, $b = 2$, and

$$y = \frac{1}{2} (e^x + e^{-x})$$

$$\frac{dy}{dx} = \frac{1}{2} (e^x - e^{-x})$$

$$\left(\frac{dy}{dx}\right)^2 = \frac{1}{4} (e^{2x} - 2 + e^{-2x})$$

$$1 + \left(\frac{dy}{dx}\right)^2 = \frac{1}{4} (e^{2x} + 2 + e^{-2x}) = \left[\frac{1}{2} (e^x + e^{-x})\right]^2.$$

Length of a Curve $y = f(x)$

The length of the curve from $x = 0$ to $x = 2$ is


$$\begin{aligned} L &= \int_0^2 \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx = \int_0^2 \frac{1}{2} (e^x + e^{-x}) dx && \text{Eq. (3) with} \\ &= \frac{1}{2} \left[e^x - e^{-x} \right]_0^2 = \frac{1}{2} (e^2 - e^{-2}) && a = 0, b = 2. \end{aligned}$$

Consider the region bounded by the graphs of $y = \ln x$, $y = 0$, and $x = e$.

- a. Find the area of the region.
- b. Find the volume of the solid formed by revolving this region about the x -axis.

$$\begin{aligned} \text{(a)} \quad A &= \int_1^e \ln x \, dx = [x \ln x]_1^e - \int_1^e dx \\ &= (e \ln e - 1 \ln 1) - [x]_1^e = e - (e - 1) = 1 \end{aligned}$$

$$\begin{aligned} \text{(b)} \quad V &= \int_1^e \pi (\ln x)^2 \, dx = \pi \left([x (\ln x)^2]_1^e - \int_1^e 2 \ln x \, dx \right) \\ &= \pi \left[\left(e (\ln e)^2 - 1 (\ln 1)^2 \right) - \left([2x \ln x]_1^e - \int_1^e 2 \, dx \right) \right] \\ &= \pi \left[e - \left((2e \ln e - 2(1) \ln 1) - [2x]_1^e \right) \right] = \pi [e - (2e - (2e - 2))] = \pi(e - 2) \end{aligned}$$



Thank you for your attention